arXiv:1703.04390v1 [math.CO] 13 Mar 2017

ON ENUMERATION OF TREE-ROOTED PLANAR CUBIC MAPS. II

YURY KOCHETKOV

Abstract. In work [4] tree-rooted planar cubic maps with marked directed edge (not in this tree) were enumerated. The number of such objects with $2n$ vertices is $C_{2n} \cdot C_{n+1}$, where C_k is Catalan number. In this work a marked directed edge is not demanded, i.e. we enumerate tree-rooted planar cubic maps. Formulas are more complex, of course, but not significantly.

1. Introduction

Plane triangulation is a planar map, where the perimeter of each face is three. The corresponding dual graph is *cubic*, i.e. the degree of each vertex is three. A plane triangulation will be called *proper*, if each edge is incident to exactly *two* faces. Otherwise it will be called *improper*.

Example 1.1.



proper triangulation



improper triangulation

The corresponding dual graphs are presented below:





A connected graph with marked directed edge will be called *edge-rooted*. Proper edge-rooted plane triangulations where enumerated by Tutte in the work [9]: the number of such triangulations with $2n$ faces is

$$\frac{2\,(4n-3)!}{n!\,(3n-1)!}.$$

A combinatorial proof of Tutte formula see in [7] (see also [1]).

All (proper and improper) plane edge-rooted triangulations were enumerated in [3] in the following way. Let F_n be the number of plane edge-rooted cubic graphs with $2n$ vertices, i.e. the number of plane edge-rooted triangulations (proper and improper) with $2n$ faces. Let us define numbers f_n, $n \geqslant -1$, in the following way:

- $f_{-1} = 1/2$;

- $f_0 = 2$;
- $f_n = (3n+2)F_n$, $n > 0$.

Then

$$f_n = \frac{4(3n+2)}{n+1} \sum_{\substack{i \geqslant -1,\ j \geqslant -1 \\ i+j=n-2}} f(i)f(j). \tag{1}$$

Thus $f_1 = 20, f_2 = 256$ and $F_1 = 4, F_2 = 32$. However, we don't know any geometric or combinatorial proof of this formula.

Example 1.2. It is easy to check the validity of this formula in cases $n = 1$ and $n = 2$. There are two cubic maps with two vertices:



Orders of their groups of automorphisms are 6 and 2, respectively.

There are six cubic maps with four vertices:



Orders of their groups of automorphisms are 1, 4, 12, 3, 2 and 2, respectively.

The list of all 26 cubic maps with six vertices see in Appendix.

Let G be a planar cubic map and T — a spanning tree. In [4] it was explained, how, using this data, to construct: a) a polygon; b) its partition into triangles by non-intersecting diagonals; c) a pairwise gluing of its edges that produces a curve of genus 0 and its triangulation.

Example 1.3. Given a cubic map with a spanning tree (thick lines and curve) we draw triangles around vertices:

After that we a) identify those edges of adjacent triangles that intersect the same edge of the spanning tree — thus we obtain a polygon; b) label identically those

sides of polygon that are intersected by the same edge of the map; c) delete the map:



If we have a polygon: a) triangulated by non-intersecting diagonals; b) its sides are pairwise identified in such way, that this identification produces a genus 0 curve, then, using these data, we can construct a planar tree-based cubic map:



In what follows by n-polygon we understand a polygon: a) with $2n$ sides; b) triangulated by non-intersecting diagonals; b) with pairwise identified sides in such way, that this identification produces a genus 0 curve. We see that the enumeration of tree-rooted planar cubic maps is equivalent to the enumeration of such objects. In work [4] we additionally mark a side of a polygon (= we mark a directed edge in a cubic map that does not belong to the spanning tree). The number of n-polygons with a marked side is $C_n \cdot C_{2n-2}$, where C_k is the Catalan number.

In this work we drop a marked side, i.e. we enumerate n-polygons (Theorem 4.1).

2. Triangulations and symmetries

Proposition 1. *Let T be a $\mathbb{Z}_k$-symmetric triangulation of a regular polygon P with $n = km$ sides, i.e. T is invariant with respect to rotations on angles $2\pi\, i/k$, $i = 1, \dots, k-1$. Then $k = 2$ or $k = 3$.*

Proof. Let R be the rotation on angle $2\pi/k$. As numbers of triangles in the triangulation T is $n-2$, then there exists a triangle $\triangle$ in T whose obtuse angle is an angle of P. As triangulation T is $\mathbb{Z}_k$-symmetric then there k copies of triangle $\triangle$ in T and their configuration is symmetric with respect to rotation R.

If we delete these triangles from polygon P, then we will obtain new polygon P_1 with $k(m-1)$ sides with triangulation T_1, which is also $\mathbb{Z}_k$-symmetric.

If we continue the process, then after $m-1$ steps we will obtain a polygon with k sides and with $\mathbb{Z}_k$-symmetric triangulation. But it is possible only when $k = 2$ or $k = 3$. □

Corollary 1. *Let P be a regular polygon with $2n$ sides, one of them — marked. Then the number of its $\mathbb{Z}_2$-symmetric triangulations is $n \cdot C_{n-1}$.*

Proof. Indeed, there are n ways to choose a big diagonal and C_{n-1} ways to triangulate a half of the polygon. □

Corollary 2. *Let P be a regular polygon with $3n$ sides, one of them — marked. Then the number of its $\mathbb{Z}_3$-symmetric triangulations is $n \cdot C_{n-1}$.*

Proof. Indeed, there are n ways to choose a $\mathbb{Z}_3$-symmetric triangle in P. The complement of this triangle consists of three symmetrically positioned polygons with $n+1$ sides each. There is C_{n-1} ways to triangulate one of them. □

3. Glueings and symmetries

Proposition 2. *Let P be a regular polygon with $2n$ sides, one of them — marked and let d_n the number of such $\mathbb{Z}_2$-symmetric pairwise glueings of its sides, that produce a genus zero curve. Then*

$$d_n = \begin{cases} \binom{2k}{k}, & \text{if } n = 2k, \\ \frac{1}{2}\binom{2k}{k}, & \text{if } n = 2k-1. \end{cases}$$

.

Proof. Let us consider a $\mathbb{Z}_2$-symmetric zero genus glueing of P. Let AB be the marked side, CD — the opposite side, KL — the glueing partner of AB and MN — the glueing partner of CD (sides KL and MN are opposite). In the counterclockwise going around of P one can meet the side KL before CD or after it. We will consider the first case and then multiply by two. The number of sides between AB and KL is even $= 2k$. We can choose any zero genus pairwise glueing of sides between AB and KL (there are C_k of them), made the symmetric glueing of sides between CD and MN and choose a $\mathbb{Z}_2$-symmetric glueing of remaining sides (there are d_{n-k-2} of them). Thus,

$$d_n = 2d_{n-2} + 2d_{n-4}C_1 + 2d_{n-6}C_2 + \ldots + 2d_2C_{(n-4)/2} + 2C_{(n-2)/2},$$

if n is even, and

$$d_n = 2d_{n-2} + 2d_{n-4}C_1 + \ldots + 2d_1C_{(n-3)/2} + C_{(n-1)/2},$$

if n is odd.

It means that

$$\left(\frac{1}{2t} + 1 + d_1t + d_2t^2 + \ldots\right) \times (2t^2 + C_1t^4 + 2C_2t^6 + \ldots) = d_1t + d_2t^2 + d_3t^3 + \ldots$$

Let

$$D(t) = \frac{1}{2t} + 1 + d_1t + d_2t^2 + \ldots$$

and let $C(t)$ be the generating function for Catalan numbers, then

$$2t^2C(t^2)D(t) = D(t) - 1 - 1/2t, \text{ or } D(t)(1 - 2t^2C(t^2)) = 1 + 1/2t.$$

As $C(t) = (1 - \sqrt{1-4t}\,)/2t$, then

$$D(t) = \left(1 + \frac{1}{2t}\right)(1-4t^2)^{-1/2} = \left(1 + \frac{1}{2t}\right)\sum_{i=0}^{\infty}\binom{2i}{i}t^{2i}. \qquad \square$$

Proposition 3. *Let P be a regular polygon with $6n$ sides, one of them — marked and let e_n the number of such $\mathbb{Z}_3$-symmetric pairwise glueings of its sides, that produce a genus zero curve. Then $e_n = \binom{2n}{n}$.*

Proof. The same reasoning gives the relation

$$2(e_{n-1} + e_{n-1}C_1 + \ldots + e_1 C_{n-2} + C_{n-1}) = e_n,$$

i.e. $2tC(t)E(t) = E(t) - 1$, where $E(t) = 1 + e_1 t + e_2 t^2 + \ldots$. Thus,

$$E(t) = \frac{1}{1 - 2tE(t)} = (1-4t)^{-1/2} = \sum_{i=0}^{\infty} \binom{2i}{i} t^i.$$

□

4. Enumeration formulas

It remains to apply Burnside formula.

Theorem 4.1. *Let t_n be the number of tree-rooted planar cubic maps with $2n$ vertices. Then*

$$(2n+2)t_n = \begin{cases} C_{n+1}C_{2n} + \frac{1}{2}(n+1)\, C_n \cdot \binom{2k}{k}, & \text{if } n+1 = 2k-1 \text{ and } n+1 \neq 3l, \\ C_{n+1}C_{2n} + (n+1)\, C_n \cdot \binom{2k}{k}, & \text{if } n+1 = 2k \text{ and } n+1 \neq 3l, \\ C_{n+1}C_{2n} + 4(2k-1)\, C_{4k-3} \cdot \binom{4k-2}{2k-1} + \frac{1}{2}(n+1)\, C_n \cdot \binom{6k-2}{3k-1}, & \\ \qquad\qquad \text{if } n+1 = 3(2k-1), & \\ C_{n+1}C_{2n} + 8k\, C_{4k-1} \cdot \binom{4k}{2k} + (n+1)\, C_n \cdot \binom{6k}{3k}, & \text{if } n+1 = 6k. \end{cases}$$

Example 4.1. If $2n = 4$ then there are 16 tree-rooted planar cubic maps. Here are they:



5. Appendix. Planar cubic maps with six vertices

We give the list of pairwise non isomorphic planar cubic maps with 6 vertices — there are 26 of them. The group of automorphisms of a map (if this group is non trivial) is also presented.



References

[1] Albenque M. and Poulalhon D., *Generic method for bijections between blossoming trees and planar maps*, arXiv: 1305.1312.

[2] Bernardi O., *Bijective counting of tree-rooted maps and shuffles of parenthesis systems*, arXiv:math/0601684.

[3] Goulden I.P. and Jackson D.M., *The KP hierarchy, branched covers, and triangulations*, arXiv: 0803.3980.

[4] Kochetkov Yu., *On enumeration of tree-rooted planar cubic maps*, arXiv: 1608.02510.

[5] Lando S. and Zvonkin A., *Graphs on surfaces and their applications. Encyclopedia of mathematical sciences* **141**, Springer-Verlag, Berlin, 2004.

[6] Mullin R.C., *On the enumeration of tree-rooted maps*, Canad. J. Math., 1967, 19, 174-183.
[7] Poulalhon D. and Schaeffer G., *Optimal coding and sampling of triangulations*, Algorithmica, 2006, 46(3), 505-527.
[8] Stanley R.P., *Enumerative combinatorics, volume 2*, Wadsworth & Brooks, 1999.
[9] Tutte W.T., *A census of planar triangulations*, Canad. J. Math., 1962, 14, 21-38.

E-mail address: yukochetkov@hse.ru, yuyukochetkov@gmail.com